

06-JUN-2005

TRILLION-P® LIQUID TURF HERBICIDE

COMMERCIAL

READ THE LABEL BEFORE USING

REGISTRATION NO: 27972
PEST CONTROL PRODUCTS ACT

GUARANTEE:

2,4-D (present as dimethyl amines)..... 190 g/L
Mecoprop-P (present as dimethyl amines)..... 100 g/L
Dicamba (present as dimethyl amines)..... 18 g/L

DO NOT FREEZE

® Registered Trademark of Plant Products Co. Ltd.

CAUTION ... POISON

Plant Products Co. Ltd.
314 Orenda Road
Brampton, Ontario
L6T 1G1

NET CONTENTS: 4L, 10L, 115L

DIRECTIONS:

Weeds Controlled: Dandelions, plantain, black medick (yellow clover), buttercup, chickweed, clover, English daisy, ground daisy, ground ivy (creeping Charlie, knotweed, ironweed), poison ivy, ragweed, shepherd's-purse, bedstraw, heal-all, devil's paint brush.

Turf (other than bentgrass): On established turf, apply at the rate of 60mL per 100m² in 10L of water. For larger areas apply at the rate of 5.5 L per hectare in 300L of water.

For bentgrass: Apply at a half rate - 30mL per 100m² in May or late August and September. Slight yellowing will occur but will disappear after about 1 week. Do not overdose bentgrass as damage or kill may occur. Make sure correct rate is applied to the whole area treated. Use 20L of water per 100m². When spraying bentgrass use only a one nozzle sprayer. Make sure spray application does not overlap or excess material will be applied to overlapped areas and damage is more likely to occur.

Maximum Application Frequency:

For good turf/lawn management, normally two applications per year per treatment site are adequate. This does not include spot treatments.

For best results apply when weeds are growing quickly, May and early June, and late August and September. On irrigated or watered areas where weeds are not under drought stress and hard to kill, apply any time. Application during drought or when weeds are under stress will result in poorer control. Do not water turf within 24 hours of application. Do not apply to newly seeded lawns. Wait until after the third mowing.

Grass seed can be safely sown 1 to 2 weeks after application at recommended rate.

Re-entry Interval for Turf:

Do not allow people (other than applicator) or pets on treatment area during application.

Do not enter treated areas until spray has thoroughly dried.

THIS PRODUCT IS NOT TO BE APPLIED BY AIR.

PRECAUTIONS:

KEEP OUT OF REACH OF CHILDREN.

Avoid contact with skin, eyes or clothing.

Protective Clothing and Equipment

Handling the concentrate (mixing and loading): wear long-sleeved shirt, long pants, coveralls or a chemical-resistant apron, chemical-resistant gloves, socks and shoes, and protective eye wear (face shield or safety glasses). Rinse gloves before removal.

Coveralls or a chemical-resistant apron should also be worn when open pouring from containers greater than 5L.

Handling the dilute spray solution (during application or repairing or cleaning equipment): wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes. Rinse gloves before removal. Gloves are not required during application when applicator is in an enclosed tractor.

Operator Use Precautions

- Wear freshly laundered clothing and clean protective equipment daily.
- Rinse gloves before removal.
- Wash hands before eating, drinking, using tobacco or using the toilet.
- If herbicide penetrates clothing remove immediately; then wash thoroughly and put on clean clothing. Throw away clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate.
- After using this product, remove clothing and launder separately, and promptly and thoroughly wash hands and exposed skin with soap and water. Follow manufacturer's instructions for cleaning personal protective clothing and equipment. If no such instructions for washables are provided, use detergent and hot water. Keep and wash personal protective equipment separately from household laundry.
- After work, remove all clothing and shower using soap and water.

Application Precautions:

Do not apply during hot, dry spells. Do not mow for at least three days before spraying.

Do not spray if rain is forecast.

Apply only when there is little or no hazard from spray drift. Small quantities of the spray, which may not be visible, may seriously injure susceptible crops such as cabbage, cauliflower, beans, turnips, tomatoes and many other broadleaf fruits, ornamentals and vegetables.

Do not use a sprayer which has contained Trillion Liquid Herbicide for spraying another pesticide on susceptible crops. Keep in original container. Do not store near seeds, fertilizers, insecticides, fungicides, feed or food.

Do not contaminate streams, lakes or ponds, irrigation water, water used for livestock or domestic purposes.

DO NOT FREEZE.

FIRST AID:

If in eyes, flush with running water for 15 minutes. If eye irritation occurs contact a physician or a poison control centre IMMEDIATELY. **If on skin**, wash thoroughly with soap and water. **If swallowed** induce vomiting (syrup of Ipecac is recommended).

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION:

Treat symptomatically.

DISPOSAL:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO THE USER:

This control product is to be used only in accordance with the directions on this label. It is an offence under the *Pest Control Products Act* to use a control product under unsafe conditions.

NOTICE TO BUYER:

Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

0405 52520

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.